

Journal of Behavioral Addictions

Serum BDNF levels in patients with gambling disorder are associated with the severity of gambling disorder and Iowa Gambling Task indices --Manuscript Draft--

Manuscript Number:	JBADD-D-15-00024R2
Full Title:	Serum BDNF levels in patients with gambling disorder are associated with the severity of gambling disorder and Iowa Gambling Task indices
Article Type:	Brief Report
Keywords:	gambling disorder; brain derived neurotrophic factor (BDNF); Iowa Gambling Task (IGT); Behavioral Addiction
Corresponding Author:	Samuel Suk-Hyun Hwang, Ph.D. Dept. of Psychology, Chonnam National University Gwangju, KOREA, REPUBLIC OF
Corresponding Author Secondary Information:	
Corresponding Author's Institution:	Dept. of Psychology, Chonnam National University
Corresponding Author's Secondary Institution:	
First Author:	Sam-Wook Choi, M.D., Ph.D.
First Author Secondary Information:	
Order of Authors:	Sam-Wook Choi, M.D., Ph.D.
	Young-Chul Shin, M.D., Ph.D.
	Jung Yeon Mok, B.A.
	Dai-Jin Kim, M.D., Ph.D.
	Jung-Seok Choi, M.D., Ph.D.
	Samuel Suk-Hyun Hwang, Ph.D.
Order of Authors Secondary Information:	
Abstract:	Background and aims: Gambling disorder (GD) shares many similarities with substance use disorders (SUDs) in clinical, neurobiological and neurocognitive features, including decision making. We evaluated the relationships among BDNF, GD and decision making, serum brain derived neurotrophic factor (BDNF) levels Methods: Twenty-one male patients with GD and 21 healthy sex- and age-matched control subjects participated. We compared the two groups in BDNF levels and examined the associations between serum BDNF levels and the Problem Gambling Severity Index (PGSI) as well as between the serum BDNF levels and Iowa Gambling Task (IGT) indices. Results: Mean serum BDNF levels were significantly increased in the patients with GD compared to healthy controls. A significant correlation between serum BDNF levels and PGSI scores was found when controlling for age, depression, and duration of GD. A significant negative correlation was obtained between serum BDNF levels and IGT improvement scores.

Serum brain-derived neurotrophic factor levels in patients with gambling disorder are associated with the severity of gambling disorder and Iowa Gambling Task indices

Abstract

Background and aims: Gambling disorder (GD) shares many similarities with substance use disorders (SUDs) in clinical, neurobiological, and neurocognitive features, including decision-making. We evaluated the relationships among, GD, decision-making, and brain-derived neurotrophic factor (BDNF), as measured by serum BDNF levels.

Methods: Twenty-one male patients with GD and 21 healthy sex- and age-matched control subjects were evaluated for associations between serum BDNF levels and the Problem Gambling Severity Index (PGSI), as well as between serum BDNF levels and Iowa Gambling Task (IGT) indices.

Results: The mean serum BDNF levels were significantly increased in patients with GD compared to healthy controls. A significant correlation between serum BDNF levels and PGSI scores was found when controlling for age, depression, and duration of GD. A significant negative correlation was obtained between serum BDNF levels and IGT improvement scores.

Discussion: These findings support the hypothesis that serum BDNF levels

constitute a dual biomarker for the neuroendocrine changes and the severity of GD in patients. Serum BDNF level may serve as an indicator of poor decision-making performance and learning processes in GD and help to identify the common physiological underpinnings between GD and SUDs.

KEY WORDS: Gambling disorder; brain-derived neurotrophic factor (BDNF); Iowa Gambling Task (IGT); behavioral addiction

1. INTRODUCTION

Gambling Disorder (GD), a type of behavioral addiction, is characterized by persistent and recurrent maladaptive gambling behavior leading to significant deleterious legal, financial, and psychosocial consequences (Grant, Kim, & Kuskowski, 2004). GD shares many similar clinical and neurobiological features with substance use disorders (SUDs), such as alterations of the mesolimbic dopamine reward pathway (Potenza, 2008), as well as neurocognitive features, including impaired decision-making.

Poor performance on the Iowa Gambling Task (IGT), designed to assess risky decision-making, has been found consistently among SUDs (Noel, Bechara, Dan, Hanak, & Verbanck, 2007). Similarly, patients with GD have demonstrated high risk-taking performance on the task (Lawrence, Luty, Bogdan, Sahakian, & Clark, 2009). Although the biological basis for decision-making is poorly understood, neural systems related to executive function and memory have been implicated (Brand,

Recknor, Grabenhorst, & Bechara, 2007).

One protein associated with various cognitive functions such as decision-making and memory is the brain-derived neurotrophic factor (BDNF) (Yamada, Mizuno, & Nabeshima, 2002). BDNF plays an important role in neuronal survival, neurogenesis, and synaptic plasticity. Studies have shown associations between BDNF and changes in behavior and psychopathology in psychiatric disorders such as depression, schizophrenia, and bipolar disorder (Monteggia et al., 2007), as well as autism spectrum disorder (Wang et al., 2015). Increases in serum levels of BDNF have been observed in drug addictions (Angelucci et al., 2010), where the involvement of BDNF in the ventral tegmental area-nucleus accumbens (VTA-NAc)-mediated processes has been implicated (Pu, Liu, & Poo, 2006).

In contrast, only a few studies have examined the association between BDNF and GD (Angelucci et al., 2013; Geisel, Banas, Hellweg, & Muller, 2012), and how BDNF levels relate to the severity of GD and the level of impairment in neurocognitive tasks remains unclear. Reduced serum BDNF has been found to relate to poor performance on the IGT (Hori, Yoshimura, Katsuki, Atake, & Nakamura, 2014) and immediate memory (Zhang et al., 2012) in patients with schizophrenia. Associations between low BDNF levels and cognitive impairment have been further confirmed in a large elderly population (Shimada et al., 2014).

In this study, we examined the relationships among GD, BDNF, and decision-making performance on the IGT in a sample of GD patients and compared the serum BDNF levels in GD patients with those in healthy control subjects. We then investigated the association of serum BDNF levels with the severity of GD and IGT indices.

2. METHODS

2.1. Participants

Twenty-one male patients who fulfilled the DSM-5 criteria for GD were recruited from the outpatient gambling clinic of the Department of Psychiatry, Gangnam Eulji Hospital, Eulji University. The diagnoses were determined by a board-certified psychiatrist (SWC) through the examination of past medical records and a semi-structured interview that included questions about the presence of co-occurring disorders. A self-report questionnaire regarding age, weight, height, alcohol-related history, regular medication use, gambling-related history, and clinical variables was also administered. The severity of GD was assessed with the Problem Gambling Severity Index (PGSI), a nine-item self-report assessment measure reported to be useful for both clinical and non-clinical settings (Young & Wohl, 2011). Mood symptoms were assessed using the Beck Depression Inventory (BDI). The exclusion criteria for the patient group were 1) any history of a chronic physical disease, 2) regular use of any medication, and 3) presence of comorbid psychiatric disorders, including alcohol and nicotine dependence. The control group consisted of 21 age- and sex-matched healthy male volunteers who had no current or past psychiatric history or history of medication use.

2.2. Measures

2.2.1. Measurement of serum BDNF levels

A total of 10 ml of blood was drawn from each subject into a serum separator tube. Samples were allowed to clot for 30 minutes before centrifugation for 15 min at

approximately 1000 g, after which the serum was removed. All samples were stored at -80°C. The serum BDNF levels were determined using an ELISA protocol according to the manufacturer's instructions (DBD00; R & D Systems, Europe).

2.2.2. IGT

For this computer-administered task, participants were asked to draw from four decks of cards. Each deck consisted of randomly distributed cards with differing amount of gains and penalties, adding up to a pre-set net outcome. Two decks contained cards with low levels of gains (e.g. \$50) and penalties (e.g. \$40), but their net outcome was favorable (e.g. \$100); the other two decks consisted of cards with high gains (e.g. \$100) but even higher penalties (e.g. \$200), so that their net outcome was unfavorable (e.g. -\$250).

All participants were instructed to try to earn as much money as possible by drawing cards one at a time from a deck of their choice. They were informed that some decks were more advantageous than others but were not told the composition of the decks. The entire IGT procedure was completed upon drawing 100 cards.

Three IGT indices were derived with high scores indicating effective strategic thinking: net total score, calculated as the number of draws from advantageous decks minus that from the disadvantageous decks (Barry & Petry, 2008); proportion of advantageous deck selections from the total number of cards; and improvement score, calculated by subtracting the net score of the first block of 20 cards from that of the last block.

2.3. Statistical analyses

An analysis of covariance, with age, body-mass index (BMI), and BDI scores entered as covariates, was used to compare serum BDNF levels of the patients and controls. The correlation between serum BDNF levels and the severity of GD based on PGSI scores in the patient group was examined using Pearson partial-correlation analysis, by controlling for age, BDI scores, and the duration of problem gambling. Finally, the association between serum BDNF levels and IGT performance was analyzed using the same method. All data are presented as means $\pm$ standard deviations (S.D.). The significance level was set at $p < 0.05$. All statistical analyses were conducted using SPSS, version 18.1 [Chicago, Illinois, USA].

2.4. Ethics

The Ethical Committee of the Eulji University, Korea, approved this study protocol. In accordance with the Declaration of Helsinki, all subjects were advised about the procedures and signed the written informed consent prior to participation.

3. RESULTS

The demographic data, gambling-related clinical variables, and IGT indices are listed in Table 1. The mean serum BDNF levels were significantly increased in the patients with GD (29051.44 ± 6237.42 pg/ml) compared to healthy controls (19279.67 ± 4375.58 pg/ml, $p < 0.0001$) (Fig. 1). We also found a significant correlation between serum BDNF levels and PGSI scores ($r = 0.56$, $p < 0.05$) after controlling for age, BDI scores, and the duration of problem gambling.

Table 1 about here

Figure 1 about here

Serum BDNF levels were also significantly negatively correlated with the IGT improvement scores ($r=-0.48$, $p<0.05$), but not with IGT total net scores ($r=-0.163$, $n.s.$) or advantageous proportion ($r=-0.19$, $n.s.$).

4. DISCUSSION

In this study, we found significantly higher serum BDNF levels among patients with GD than in healthy controls, as well as a positive association between the serum BDNF levels and severity of GD. Such findings are in partial agreement with previous studies showing that serum BDNF levels increased in GD (Angelucci et al., 2013; Geisel et al., 2012), although these studies present different results regarding the association between serum BDNF levels and severity of GD. Such discrepancies might be related to external factors that affect serum BDNF levels, including BMI, depression, and other confounding factors (Piccinni et al., 2008). Together with these two previous studies (Angelucci et al., 2013, Geisel et al., 2012), our findings suggest that behavioral addictions might be associated with neural plasticity similar to alterations observed in SUDs. Increased serum BDNF levels might then represent a compensatory mechanism to normalize dopaminergic transmission in VTA and the

1 169 NAc (Geisel et al., 2012). Another plausible explanation is that increased BDNF
2
3 170 plays a role in neuroprotective and stress-preventive processes in patients with GD,
4
5 171 especially during stressful situations, as found in those with SUDs (Bhang, Choi, &
6
7
8 172 Ahn, 2010; Geisel et al., 2012).

10 173 Although a recent study (Kang et al., 2010) showed that the BDNF Val66Met
11
12 174 polymorphism may affect decision-making performance as measured by IGT, to the
13
14 175 best of our knowledge, our study is the first to demonstrate a significant association
15
16 176 between serum BDNF levels and IGT improvement scores. The IGT improvement
17
18 177 score particularly reflects learning processes based on the evaluation of choice-
19
20 178 outcomes of rewards and penalties leading to long-term gain or loss. This learning
21
22 179 involves discounting immediate rewards while formulating an advantageous strategy
23
24 180 based on previous cumulative outcomes. A recent study (Kräplin et al., 2014) found
25
26 181 problem gamblers showed higher overall impulsivity compared to healthy controls
27
28 182 and higher 'choice impulsivity' compared to a Tourette syndrome group, but similar
29
30 183 levels of impulsivity as an alcohol-dependent group. Higher BDNF concentration has
31
32 184 also been positively correlated with higher impulsivity in PTSD patients (Martinotti et
33
34 185 al., 2015) suggesting that impulsiveness may be associated with greater BDNF
35
36 186 expression. Additionally, in murine models, BDNF has been implicated in the actions
37
38 187 of serotonergic neurons, particularly in aggression and impulsivity (Lyons et al.,
39
40 188 1999). Both BDNF and serotonin regulate the development and plasticity of neural
41
42 189 circuits in mood disorders (Martinowich & Lu, 2008). In humans, BDNF Val66Met
43
44 190 polymorphism in schizophrenia patients has been associated with aggressive
45
46 191 behavior (Spalletta et al., 2010), while serotonin has been found to play a significant
47
48 192 role in learning and memory (Meneses & Liy-Salmeron, 2012). Taken together, our
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 193 results suggest that BDNF may also play a role in learning processes, and that the
2
3 194 relationship between BDNF and serotonin needs to be further examined.
4

5
6 195 Some limitations of this study warrant discussion; our sample size was modest
7
8 196 and contained only male GD patients, thus limiting the generalizability of our results.
9
10 197 Serum BDNF levels were examined rather than central nervous system BDNF
11
12 198 levels. Although BDNF regulation in peripheral blood is still poorly understood,
13
14 199 peripheral concentrations are widely used as a mirror of the same brain parameter
15
16 200 (Yamada, Mizuno, & Nabeshima, 2002). Because BDNF is known to cross the blood-
17
18 201 brain barrier in both directions, a substantial part of peripheral BDNF might originate
19
20 202 from neuronal cells of the central nervous system (Karege, Schwald, & Cisse, 2002).
21
22 203 At the present, the relationships among BDNF, severity of disorder, and decision-
23
24 204 making in GD patients are not clearly delineated, and future studies should consider
25
26 205 these limitations in their designs for a better understanding of such relationships. In
27
28 206 addition, we did not take into account personality factors in our study design.
29
30 207 Previous studies have suggested relationships between pathological gambling and
31
32 208 personality characteristics such as novelty-seeking and self-directedness (Jiménez-
33
34 209 Murcia et al., 2010; Martinotti et al., 2006), but consensus on the relationship
35
36 210 between the BDNF levels and these personality traits has yet to be reached due to
37
38 211 inconsistent results (Maclaren, Fugelsang, Harrigan, & Dixon, 2011). The results of
39
40 212 our study should be interpreted carefully in light of such a limitation.
41
42
43
44
45
46
47
48
49
50
51

52 214 **5. CONCLUSION**

53
54
55

56 215 The findings of this study support the hypothesis that serum BDNF levels may serve
57
58 216 as a candidate biomarker for neural plasticity and the severity of GD in these
59
60
61
62
63
64
65

patients. Furthermore, heightened serum BDNF levels in GD may indicate poor decision-making performance, a characteristic feature of SUDs. This study therefore is a meaningful addition to the growing body of research supporting the common neurobiological underpinnings of SUDs and GD.

References

- Angelucci, F., Martinotti, G., Gelfo, F., Righino, E., Conte, G., Caltagirone, C., Bria, P., & Ricci, V. (2013). Enhanced BDNF serum levels in patients with severe pathological gambling. *Addiction Biology*, 18, 749-751.
- Angelucci, F., Ricci, V., Martinotti, G., Palladino, I., Spalletta, G., Caltagirone, C., & Bria, P. (2010). Ecstasy (MDMA)-addicted subjects show increased serum levels of brain-derived neurotrophic factor, independently from a rise of drug-induced psychotic symptoms. *Addiction Biology*, 15, 365-367.
- Barry, D., & Petry, N. M. (2008). Predictors of decision-making on the Iowa Gambling Task: independent effects of lifetime history of substance use disorders and performance on the Trail Making Test. *Brain and Cognition*, 66, 243-252.
- Bhang, S. Y., Choi, S. W., & Ahn, J. H. (2010). Changes in plasma brain-derived neurotrophic factor levels in smokers after smoking cessation. *Neuroscience Letters*, 468, 7-11.
- Brand, M., Recknor, E. C., Grabenhorst, F., & Bechara, A. (2007). Decisions under ambiguity and decisions under risk: correlations with executive functions and comparisons of two different gambling tasks with implicit and explicit rules. *Journal of Clinical and Experimental Neuropsychology*, 29, 86-99.
- Geisel, O., Banas, R., Hellweg, R., & Muller, C. A. (2012). Altered serum levels of

- brain-derived neurotrophic factor in patients with pathological gambling.
European Addiction Research, 18, 297-301.
- Grant, J. E., Kim, S. W., & Kuskowski, M. (2004). Retrospective review of treatment retention in pathological gambling. *Comprehensive Psychiatry*, 45, 83-87.
- Hori, H., Yoshimura, R., Katsuki, A., Atake, K., & Nakamura, J. (2014). Relationships between brain-derived neurotrophic factor, clinical symptoms, and decision-making in chronic schizophrenia: data from the Iowa Gambling Task. *Frontiers of Behavioral Neuroscience*. 8, 417. doi: 10.3389/fnbeh.2014.00417
- Jiménez-Murcia, S., Alvarez-Moya, E. M., Stinchfield, R., Fernández-Aranda, F., Granero, R., Aymamí, N., Gómez-Peña, M., Jaurieta, N., Bove, F., & Menchón, J. M. (2010). Age of onset in pathological gambling: clinical, therapeutic and personality correlates. *Journal of Gambling Studies*. 26, 235-248.
- Kang, J. I., Namkoong, K., Ha, R. Y., Jhung, K., Kim, Y. T., & Kim, S. J. (2010). Influence of BDNF and COMT polymorphisms on emotional decision making. *Neuropharmacology*, 58, 1109-1113.
- Karege, F., Schwald, M., & Cisse, M. (2002). Postnatal developmental profile of brain-derived neurotrophic factor in rat brain and platelets. *Neuroscience Letters*, 328, 261-264.
- Kräplin, A., Bühringer, G., Oosterlaan, J., van den Brink, W., Goschke, T., & Goudriaan, A. E. (2014). Dimensions and disorder specificity of impulsivity in pathological gambling. *Addictive Behavior*, S0306-4603(14), 00170-1. doi: 10.1016/j.addbeh.2014.05.021.
- Lawrence, A. J., Luty, J., Bogdan, N. A., Sahakian, B. J., & Clark, L. (2009). Impulsivity and response inhibition in alcohol dependence and problem gambling. *Psychopharmacology*, 207, 163-172.
- Lyons, W. E., Mamounas, L. A., Ricaurte, G. A., Coppola, V., Reid, S. W., Bora, S. H., Wihler, C., Koliatsos, V. E., & Tessarollo, L. (1999). Brain-derived neurotrophic

- factor-deficient mice develop aggressiveness and hyperphagia in conjunction with brain serotonergic abnormalities. *Proceedings of the National Academy of Sciences*, 96, 15239-15244.
- Maclaren, V. V., Fugelsang, J. A., Harrigan, K. A., & Dixon, M. J. (2011). The personality of pathological gamblers: a meta-analysis. *Clinical Psychology Review*, 31, 1057-1067.
- Martinotti, G., Andreoli, S., Giametta, E., Poli, V., Bria, P., & Janiri, L. (2006). The dimensional assessment of personality in pathologic and social gamblers: the role of novelty seeking and self-transcendence. *Comprehensive Psychiatry*, 47(5), 350-356;
- Martinotti, G., Sepede, G., Brunetti, M., Ricci, V., Gambi, F., Chillemi, E., Vellante, F., Signorelli, M., Pettorruso, M., De Risio, L., Aguglia, E., Angelucci, F., Caltagirone, C., & Di Giannantonio, M. (2015). BDNF concentration and impulsiveness level in post-traumatic stress disorder. *Psychiatry Research*, 229, 814-818.
- Martinowich, K., & Lu, B. (2008). Interaction between BDNF and serotonin: role in mood disorders. *Neuropsychopharmacology*, 33, 73-83.
- Meneses, A., & Liy-Salmeron, G. (2012). Serotonin and emotion, learning and memory. *Review of Neuroscience*, 23, 543-553.
- Monteggia, L., Lukiart, B., Barrot, M., Theobald, D., Malkovska, I., Nef, S., Parada, L. F., & Nestler, E. J. (2007). Brain-derived neurotrophic factor conditional knockouts show gender differences in depression related behaviours. *Biological Psychiatry*, 61, 187-197.
- Noel, X., Bechara, A., Dan, B., Hanak, C., Verbanck, P. (2007). Response inhibition deficit is involved in poor decision making under risk in nonamnesic individuals with alcoholism. *Neuropsychology*, 21, 778-786.
- Piccinni, A., Marazziti, D., Del Debbio, A., Bianchi, C., Roncaglia, I., Mannari, C., Origlia, N., Catena, D. M., Massimetti, G., Domenici, L., & Dell'Osso, L. (2008).

- Diurnal variation of plasma brain-derived neurotrophic factor (BDNF) in humans: an analysis of sex differences. *Chronobiology International*, 25, 819-826.
- Potenza, M. N. (2008). Review: The neurobiology of pathological gambling and drug addiction: an overview and new findings. *Philosophical transactions of the Royal Society of London Series B, Biological Sciences*, 363, 3181-3189.
- Pu, L., Liu, Q. S., & Poo, M. M. (2006). BDNF-dependent synaptic sensitization in midbrain dopamine neurons after cocaine withdrawal. *Nature Neuroscience*, 9, 605-607.
- Shimada, H., Makizako, H., Doi, T., Yoshida, D., Tsutsumimoto, K., Anan, Y., Uemura, K., Lee, S., Park, H., & Suzuki, T. (2014). A large, cross-sectional observational study of serum BDNF, cognitive function, and mild cognitive impairment in the elderly. *Frontiers in Aging Neuroscience*, 15, 6:69. doi: 10.3389/fnagi.2014.00069.
- Spalletta, G., Morris, D. W., Angelucci, F., Rubino, I. A., Spoletini, I., Bria, P., Martinotti, G., Siracusano, A., Bonaviri, G., Bernardini, S., Caltagirone, C., Bossù, P., Donohoe, G., Gill, M., & Corvin, A. P. (2010). BDNF Val66Met polymorphism is associated with aggressive behavior in schizophrenia. *European Psychiatry*, 25, 311-313.
- Wang, M., Chen, H., Yu, T., Cui, G., Jiao, A., & Liang, H. (2015). Increased serum levels of brain-derived neurotrophic factor in autism spectrum disorder. *Neuroreport*, 26, 638-641.
- Yamada, K., Mizuno, M., & Nabeshima, T. (2002). Role for brain-derived neurotrophic factor in learning and memory. *Life Science*, 70, 735-744.
- Young, M. M., & Wohl, M. J. (2011). The Canadian Problem Gambling Index: an evaluation of the scale and its accompanying profiler software in a clinical setting. *Journal of gambling studies / co-sponsored by the National Council on Problem Gambling and Institute for the Study of Gambling and Commercial*

Gaming, 27, 467-485.

Zhang, X. Y., Liang, J., Chen da, C., Xiu, M. H., Yang, F.D., Kosten, T. A., & Kosten, T. R. (2012). Low BDNF is associated with cognitive impairment in chronic patients with schizophrenia. *Psychopharmacology (Berl)*, 222(2), 277-284.

Figure Legends

Figure 1. The mean serum BDNF levels were significantly increased in the patients with gambling disorder (29051.44 ± 6237.42 pg/ml) compared to healthy controls (19279.67 ± 4375.58 pg/ml, $p < 0.0001$) by ANCOVA with age, BMI, and the scores of BDI as covariates. The box plots show the median and quartiles, and the whisker caps of the box plots show the mean 5th and 95th percentile values.

* Indicates statistical significance ($F=12.11$, $p < 0.001$).




Table 1 Demographic data, BDI, BDNF, IGT index, and GD related variables

Variable	GD (<i>n</i> =21)	Control (<i>n</i> =21)	Test statistics	<i>p</i> -value
	<i>M</i> (<i>SD</i>)	<i>M</i> (<i>SD</i>)		
Age	40.52 (12.35)	39.29 (3.96)	<i>t</i> = 0.438	0.664
BMI	25.17 (3.42)	22.54 (2.43)	<i>t</i> = 2.873	<0.01
BDI	18.48 (11.78)	4.10 (3.03)	<i>t</i> = 5.420	<0.0001
BDNF(pg/ml)	29051.44(6237.42)	19279.67(4375.58)	<i>t</i> = 5.877	<0.0001
IGT total Net score	9.14 (21.81)			
Advantageous proportion	0.55 (0.11)			
IGT improvement score	2.86 (5.08)			
CPGI-PGSI	20.10 (4.79)			
GD duration (years)	8.14 (5.30)			
No. of gambling methods			$\chi^2= 0.048$	0.827
One	10 (47.6%)			
Multiple(two or more)	11 (52.4%)			
GD type			$\chi^2= 2.333$	0.127
Action type	14 (66.7%)			
Escape type	7 (33.3%)			
Gambling category ^a			$\chi^2= 2.333$	0.127
Strategic	7 (33.3%)			
Analytic	14 (66.7%)			

Note. * Marked variables are categorical variables with N (%), therefore Chi-square test was used. GD: Gambling Disorder, BMI: Body Mass Index (weight / height²), BDI: Beck Depression Inventory, BDNF: Brain-derived neurotrophic factor, IGT total Net score: total advantageous deck counts minus total disadvantageous deck counts, Advantageous proportion: advantageous deck counts / total card selection (100 cards), IGT improvement score: block5 IGT Net score minus block1 IGT Net score, CPGI-PGSI: Canadian Problem Gambling Index-Problem Gambling Severity Index, ^a Strategic: Casino gambling (e.g. Black-Jack); Analytic: Sports betting, Horse racing, Bicycle racing, Motor boat racing, Stock-trading